arXiv:1202.0985v1 [math.CV] 5 Feb 2012

LÊ'S POLYHEDRON FOR LINE SINGULARITIES

AURÉLIO MENEGON NETO

Abstract. We study the topology of a line singularity, which is a complex hypersurface with non-isolated singularity given by a smooth curve, by means of the description of the degeneration of its Milnor fibre to the singular variety. We construct a polyhedron P_t in $\mathring{X}_t$, the interior of the Milnor fibre, a polyhedron P_0 in $\mathring{X}_0$, the interior of the singular fibre, and a continuous map taking $\mathring{X}_t$ to $\mathring{X}_0$ and P_t to P_0, which restricts to a homeomorphism outside these polyhedra.

1. Introduction

The idea of studying the critical level of a complex function by studying the non-critical level is classical, used by many authors like Milnor, Hirzebruch, Brieskorn, Pham and others. This lead to the classic Fibration Theorem of Milnor and to the study of the vanishing homology of a singularity.

In the case of an isolated singularity, Lê Dung Trang refined the idea of vanishing homology and proved that there exists a polyhedron in the Milnor fibre such that the Milnor fibre deformation retracts to such polyhedron, and that there is a continuous map from the Milnor fibre to the singular one which restricts to a homeomorphism outside the polyhedron and takes the polyhedron to the singular point [7].

It is unlikely that there is a natural extension of this result to holomorphic functions with arbitrary singular set. In this paper we show that there is such theorem for an important class of singularities called *line singularities*, defined and first studied by D. Siersma in [11]. These are nothing but complex singularity germs with singular set a complex smooth curve.

Line singularities are closely related to the μ-constant family problem, since in many cases one can see a family of complex isolated singularity hypersurfaces depending holomorphically on one parameter as a line singularity. In fact, if f is a line singularity defined by a family of isolated singularities f_s, then the Lê Polyhedron P_t of f describes the degeneration of the Milnor fibre of f_s to the Milnor fibre of f_0.

In particular, in the last section of the present work we prove, by means of the Lê Polyhedra, that if such a family is μ-constant and admits an uniform Milnor radius, then it must be equisingular. Although this result is an immediate consequence of some classical theorems of K. Bekka about the (c)-regularity of a family of isolated singularity hypersurfaces ([1], [2], [3]), the proof presented here provides a new point of view into this class of problems.

Date: 29-01-2012.

The author thanks for the financial support received from CONACYT, Mexico, from the "Laboratorio Internacional Solomon Lefschetz" (LAISLA), associated to the CNRS (Francia) and CONACYT (Mexico) and from the "Instituto de Matemáticas de Cuernavaca".

2. Main theorem and structure of the proof

Consider $f : (\mathbb{C}^{n+1}, 0) \to (\mathbb{C}, 0)$ a line singularity with singular set a the smooth complex curve Σ and for any positive real ϵ, denote by $\mathbf{B}_\epsilon$ the ball in $\mathbb{C}^{n+1}$ centered at 0 and with radius ϵ.

We say that $\epsilon > 0$ is an uniform Milnor radius for f and that $\mathbf{B}_\epsilon$ is a uniform Milnor ball for f if, for any $s \in \Sigma$, the intersection $\mathbf{B}_\epsilon \cap H_s$ is a Milnor ball for the restriction of f to H_s, where H_s is a family of hyperplane sections of $\mathbb{C}^n$ transversal to Σ at s.

Also denote by $\mathring{X}_t$ the interior of X_t. We show that any line singularity admits a Lê Polyhedron in the following sense:

Theorem 1. *If $f : (\mathbb{C}^{n+1}, 0) \to (\mathbb{C}, 0)$ is a line singularity, then there exist ϵ and η sufficiently small, with $0 < \eta << \epsilon << 1$, such that for any $t \in \boldsymbol{D}^*_\eta$:*

- (i) *There exists a polyhedron P_t of real dimension $n+1$ in the Milnor fibre X_t such that X_t deformation retracts to P_t;*
- (ii) *If f does not admit a uniform Milnor radius, there exist a retractable polyhedron P_0 of real dimension $n+1$ in the singular fibre X_0 such that X_0 deformation retracts to P_0; and there is a continuous map $\Psi_t : \mathring{X}_t \to \mathring{X}_0$ which sends P_t to P_0 and such that Ψ_t restricts to a homeomorphism from $\mathring{X}_t \backslash P_t$ to $\mathring{X}_0 \backslash P_0$;*
- (iii) *If f admits a uniform Milnor radius, there exists collapsing map $\Psi_t : \mathring{X}_t \to \mathring{X}_0$ sending P_t to $\Sigma \cap \boldsymbol{B}_\epsilon$ and such that Ψ_t restricts to a homeomorphism from $\mathring{X}_t \backslash P_t$ to $\mathring{X}_0 \backslash (\Sigma \cap \boldsymbol{B}_\epsilon)$.*

By polyhedron we mean a topological space which admits a triangulation by a simplicial complex (see [12]).

The proof of this theorem is presented in section 3, and it is based on lemmas 4, 5, 6 and 7. Lemmas 4 and 5 are particular cases of propositions 8 and 9, respectively, presented in section 4, and which concern the construction of a Lê Polyhedron for complex analytic singularities $f : M \to \mathbb{C}$ defined on a compact n-dimensional complex manifold with boundary ∂M such that the singular set of f is the whole special fibre $M_0 := f^{-1}(0)$. Here, by a compact complex manifold with boundary we mean that M is compact, smooth and its interior is a complex manifold.

We then prove propositions 8 and 9 in section 5, when $n = 2$, and in section 6, when $n \geq 3$. Although the ideas of the proofs of these propositions are quite analogous to those presented in [7], there are mainly two significative differences: First, in our case the spaces M, M_t and M_0 are smooth manifolds, so we do not have to deal with stratifications here. Second, our study is global, that is, we shall look at all the space M.

Remark 2. *Following exactly the same steps of sections 4, 5 and 6, it is an easy exercise to construct a Lê Polyhedron for an isolated singularity $f : \boldsymbol{B}_\epsilon \to \mathbb{C}$ such that $0 \in \boldsymbol{B}_\epsilon$ is the only singularity of f and such that $f^{-1}(0)$ intersects $\boldsymbol{S}_\epsilon := \partial \boldsymbol{B}_\epsilon$ transversally, without the assumption that ϵ is a Milnor radius for f, that is, in a global sense. In other words, one can show that for any t sufficiently small such that $f^{-1}(t)$ intersects $\boldsymbol{S}_\epsilon$ transversally (including $t = 0$), there exists a polyhedron P_t in $f^{-1}(t)$ such that $f^{-1}(t)$ deformation retracts to P_t (note that $f^{-1}(0)$ is not necessarily contractible). Moreover, for any t small there exists a continuous map $\Psi_t : f^{-1}(t) \to f^{-1}(0)$ which restricts to a homeomorphism from $f^{-1}(t) \backslash P_t$ to*

$f^{-1}(0)\backslash P_0$ and sends P_t to P_0. Clearly, $P_0 = \{0\}$ if ϵ is a Milnor radius for f. This is going to be useful in the prove of theorem 1, in the next section.

Finally, in section 7 we prove lemmas 4, 5, 6 and 7.

The author would like to thank J. Seade, who introduced him to the subjects concerned in this work and made significant contributions to it.

---

3. Functions with singular set a line

Let $f : (\mathbb{C}^{n+1}, 0) \to (\mathbb{C}, 0)$ be a line singularity, that is, a holomorphic germ of a function whose singular set Σ is a smooth complex curve. Without lost of generality, we can suppose that Σ is the z_n-axis, that is,

$$\Sigma = \{z_1 = \cdots = z_n = 0\},$$

and for any $s \in \Sigma$, consider the hyperplane section H_s given by

$$H_s := \{z_{n+1} = s\}.$$

Then considering the restrictions

$$f_s := f_{|H_s} : \mathbb{C}^n \to \mathbb{C}$$

we obtain a family, in the parameter $s \in \Sigma \overset{\text{diff.}}{=} \mathbb{C}$, of holomorphic functions with isolated singularity.

It is well known that one can define the Milnor fibration of f using polydiscs, that is, if ϵ_1 and ϵ_2 are sufficiently small positive reals with $0 < \epsilon_1 < \epsilon_2 << 1$, one has that the restriction

$$f_| : f^{-1}(\mathbf{D}^*_\eta) \cap (\mathbf{B}_{\epsilon_1} \times \mathbf{B}_{\epsilon_2}) \to \mathbf{D}^*_\eta$$

is a fibre bundle and $f^{-1}(t) \cap \mathbf{B}_\epsilon$ is homeomorphic to $f^{-1}(t) \cap (\mathbf{B}_{\epsilon_1} \times \mathbf{B}_{\epsilon_2})$, where

- $\mathbf{D}_\eta$ denotes the disk in $\mathbb{C}$ centered at zero and with radius η;
- $\mathbf{D}^*_\eta$ is the punctured disk $\mathbf{D}_\eta \backslash \{0\}$;
- $\mathbf{B}_{\epsilon_1}$ denotes the ball in $\Sigma \overset{\text{diff.}}{=} \mathbb{C}$ centered at zero and with radius ϵ_1;
- $\mathbf{B}_{\epsilon_2}$ denotes the ball in $\mathbb{C}^n$ centered at zero and with radius ϵ_2;
- $\mathbf{B}_\epsilon \subset \mathbb{C}^{n+1}$ is an usual Milnor ball for f.

Now, since either Σ or $\Sigma\backslash\{0\}$ is a stratum of a Whitney stratification for f, we can take ϵ_1 and ϵ_2 sufficiently small such that:

- $\big(f^{-1}(0)\backslash f_0^{-1}(0)\big) \cap (\mathbf{B}_{\epsilon_1} \times \mathbf{B}_{\epsilon_2})$ is a fibre bundle over $(\Sigma \cap \mathbf{B}_{\epsilon_1})\backslash\{0\}$ with fibre $f_s^{-1}(0) \cap (\{s\} \times \mathbf{B}_{\epsilon_2})$;
- $(\{0\} \times \mathbf{B}_{\epsilon_2})$ is a Milnor ball for f_0.

Then for any $t \in \mathbf{D}^*_\eta$, let X_t denote the Milnor fibre of f, that is,

$$X_t := f^{-1}(t) \cap (\mathbf{B}_{\epsilon_1} \times \mathbf{B}_{\epsilon_2}),$$

which is a compact smooth manifold of dimension n, and let X_0 denote the singular fibre of f, that is,

$$X_0 := f^{-1}(0) \cap (\mathbf{B}_{\epsilon_1} \times \mathbf{B}_{\epsilon_2}),$$

which is a compact singular hypersurface of dimension n whose singular set is

$$\Sigma_\epsilon := \Sigma \cap \mathbf{B}_{\epsilon_1}.$$

Also, to simplify notation, from now on we shall denote by f_s the restriction of f to the intersection $H_s \cap (\{s\} \times \mathbf{B}_{\epsilon_2})$.

Now, for any $t \in \mathbf{D}_\eta^*$, let

$$\pi_t : \mathring{X}_t \to \Sigma_\epsilon$$

be the analytic function given by the natural projection of $\mathbb{C}^{n+1}$ on Σ restricted to $\mathring{X}_t$. Considering ϵ_1 and ϵ_2 sufficiently small, there are two possibilities:

- (a) Σ_ϵ is a stratum of a Whitney stratification for f; then π_t is a (trivial) topological fibre bundle over Σ_ϵ and therefore $0 \in \mathbb{C}$ is a regular value for π_t;
- (b) $\Sigma_\epsilon \backslash \{0\}$ is a stratum of a Whitney stratification for f; then π_t is a topological fibre bundle over $\Sigma_\epsilon \backslash \{0\}$ and $0 \in \mathbb{C}$ is a critical value for π_t; there are two cases to consider:
  - (b_1) $Sing(\pi_t) \subsetneq \pi_t^{-1}(0)$ and therefore $\pi_t^{-1}(0)$ is singular;
  - (b_2) $Sing(\pi_t) = \pi_t^{-1}(0)$ and therefore $\pi_t^{-1}(0)$ is smooth.

Note that, for any $t \in \mathbf{D}_\eta^*$ and $s \in \Sigma_\epsilon$, we have

$$\pi_t^{-1}(s) = int[f_s^{-1}(t)]$$

But then $\pi_t^{-1}(0) = int[f_0^{-1}(t)]$, the Milnor fibre of the isolated singularity function f_0, and therefore it is smooth. So case (b_1) cannot occur.

3.1. **Case** (a)**.** In this case, if $\epsilon_2 > 0$ is sufficiently small, clearly $\mathbf{B}_{\epsilon_2}$ is a Milnor ball for any any f_s with $s \in \Sigma_\epsilon$, and then we can construct a Lê Polyhedron $P_t \subset \mathring{X}_t$ as follows:

For each $s \in \Sigma_\epsilon$, let $P_{t,s} \subset f_s^{-1}(t)$ be a Lê Polyhedron for the isolated singularity function f_s as in [7]. Then the fibre bundle

$$\begin{tikzcd} int[f_s^{-1}(t)] \arrow[r, hook] & \mathring{X}_t \arrow[d, "\pi_t"] \\ & \Sigma_\epsilon \end{tikzcd}$$

induces a fibre bundle

$$\begin{tikzcd} P_{t,s} \arrow[r, hook] & P_t \arrow[d] \\ & \Sigma_\epsilon \end{tikzcd}$$

where the total space P_t is a *Lê Polyhedron for* f.

In fact, since $f_s^{-1}(t)$ deformation retracts to $P_{t,s}$, it follows that X_t deformation retracts to P_t. Moreover, we know that for each $s \in \Sigma_\epsilon$, there exists a collapsing map

$$\Psi_{t,s} : f_s^{-1}(t) \to f_s^{-1}(0)$$

which restricts to a homeomorphism $f_s^{-1}(t) \backslash P_{t,s} \to f_s^{-1}(0) \backslash \{s\}$ and such that $\Psi_{t,s}(P_{t,s}) = \{s\}$. Then we can define a continuous collapsing map

$$\Psi_t : \mathring{X}_t \to \mathring{X}_0$$

for f setting

$$\Psi_t(z) := \Psi_{t,\pi_t(z)}(z).$$

It is obvious that Ψ_t restricts to a homeomorphism $\overset{\circ}{X}_t \backslash P_t \to \overset{\circ}{X}_0 \backslash \Sigma_\epsilon$ and that $\Psi_t(P_t) = \Sigma_\epsilon$.

Remark 3. Note that P_t obviously deformation retracts to $P_{t,0}$, which has the homotopy type of a bouquet of $\mu(f_0)$-spheres of dimension $(n-1)$, as we shall see later, in lemma 4.

3.2. **Case** (b_2)**.** In this case we have to understand how $int[f_s^{-1}(t)] = \pi_t^{-1}(s)$ degenerates to $int[f_0^{-1}(t)] = \pi_t^{-1}(0)$ as $s \in \Sigma_\epsilon \backslash \{0\}$ goes to $0 \in \Sigma_\epsilon$, for any $t \in \mathbf{D}_\eta^*$ fixed. To do that, we use lemmas 4 and 5 bellow, which give a Lê Polyhedron for the projection π_t, and lemmas 6 and 7. We shall prove these lemmas in the next sections.

Lemma 4. *For any $t \in \boldsymbol{D}_\eta^*$ and $s \in \Sigma_\epsilon$, one has that:*

- (i) *there exists a polyhedron $P_{t,s}$ in $\pi_t^{-1}(s)$, of real dimension $n-1$, which has the homotopy type of a bouquet of spheres of dimension $n-1$, that is,*
$$P_{t,s} \overset{\text{homot.}}{\simeq} \bigvee_{\mu_s} \boldsymbol{S}^{n-1},$$
where $\mu_s \geq 0$ is the number of spheres, with μ_0 being the Milnor number of the isolated singularity f_0 and $\mu_s \geq \mu_0$;
- (ii) *$\pi_t^{-1}(s)$ deformation retracts to $P_{t,s}$.*

Lemma 5. *For any $t \in \boldsymbol{D}_\eta^*$ and $s \in \Sigma_\epsilon \backslash \{0\}$, there exists a continuous map $\tilde{\Psi}_{t,s} : \pi_t^{-1}(s) \to \pi_t^{-1}(0)$ such that $\tilde{\Psi}_{t,s}$ restricts to a homeomorphism $\pi_t^{-1}(s) \backslash P_{t,s} \to \pi_t^{-1}(0) \backslash P_{t,0}$ and takes $P_{t,s}$ to $P_{t,0}$.*

Lemma 6. *The construction of the polyhedra $P_{t,s}$ and of the collapses $\tilde{\Psi}_{t,s} : \pi_t^{-1}(s) \to \pi_t^{-1}(0)$ can be done simultaneously for any $s \in \Sigma_\epsilon \backslash \{0\}$. Then we obtain a polyhedron P_t in X_t, of real dimension $n+1$, such that X_t deformation retracts to P_t and such that for any $s \in \Sigma_\epsilon$, the Milnor fibre $\pi_t^{-1}(s)$ deformation retracts to the intersection $P_t \cap \pi_t^{-1}(s)$.*

Lemma 7. *For any $t \in \boldsymbol{D}_\eta^*$ and $s \in \Sigma_\epsilon$, the polyhedron $P_{t,s}$ is also a Lê Polyhedron for f_s, in the global sense* (see remark 2).

Then the polyhedron P_t is a Lê Polyhedron for f, since we can define a collapsing map
$$\Psi_t : \overset{\circ}{X}_t \to \overset{\circ}{X}_0$$
for f setting
$$\Psi_t(z) := \Psi_{t,\pi_t(z)}(z),$$
where $\Psi_{t,s} : f_s^{-1}(t) \to f_s^{-1}(0)$ is a collapsing map for f_s (in the globally sense), which restricts to a homeomorphism from $f_s^{-1}(t) \backslash P_{t,s}$ to $f_s^{-1}(0) \backslash P_{0,s}$ and sends $P_{t,s}$ to $P_{0,s}$ (where $P_{0,s}$ is a special polyhedron for f_s in the global sense, see remark 2). Set
$$P_0 := \bigcup_{s \in \Sigma_\epsilon} P_{0,s},$$
a polyhedron in X_0 containing Σ_ϵ and such that X_0 deformation retracts to P_0 (and hence P_0 obviously deformation retracts to $\{0\}$, since X_0 does). Then Ψ_t clearly

restricts to a homeomorphism $\mathring{X}_t \backslash P_t \to \mathring{X}_0 \backslash P_0$ and $\Psi_t(P_t) = P_0$, and then we have proved (ii) of theorem 1.

Now suppose that f admits an uniform Milnor radius. Then f_s is an isolated singularity and we can consider $\Psi_{t,s} : int[f_s^{-1}(t)] \to int[f_s^{-1}(0)]$ to be a collapsing map for f_s (in the local sense of [7]), which restricts to a homeomorphism from $int[f_s^{-1}(t)] \backslash P_{t,s}$ to $int[f_s^{-1}(0)] \backslash \{s\}$ and sends $P_{t,s}$ to $\{s\}$. Hence Ψ_t restricts to a homeomorphism $\mathring{X}_t \backslash P_t \to \mathring{X}_0 \backslash \Sigma_\epsilon$ and $\Psi_t(P_t) = \Sigma_\epsilon$, so we have proved (iii) of Theorem 1.

4. The Lê polyhedron for a special class of singularities

Let M be a compact oriented smooth n-dimensional complex manifold in $\mathbb{C}^{n+1}$, with boundary ∂M, and let $f : M \to \mathbb{C}$ be a holomorphic function such that:

(i) the critical set of f is given by $\Sigma = f^{-1}(0) \neq \emptyset$;

(ii) $f^{-1}(t)$ is a connected hypersurface in M which intersects ∂M transversally, for any $t \in \mathbb{C}$ sufficiently small.

Later we will take M to be the Milnor fibre of a complex hypersurface. For any positive real $\eta > 0$, define

$$M_\eta := f^{-1}(\mathbf{D}_\eta).$$

By Ehresmann's fibration lemma, the restriction

$$f_| : M_\eta \backslash (f^{-1}(0) \cap M_\eta) \to \mathbf{D}^*_\eta$$

is a topological fibre bundle. We want to describe how $M_t := f^{-1}(t)$ degenerates to $M_0 := f^{-1}(0)$, as $t \in \mathbb{C}$ goes to zero. We have the following propositions, which generalize lemmas 4 and 5 of the previous section:

Proposition 8. *For any $t \in \boldsymbol{D}_\eta$, one has that:*

(i) *there exists a polyhedron P_t in M_t, of real dimension $n-1$, which has the homotopy type of a bouquet of spheres of (distinct) dimensions $\leq n-1$, that is,*

$$P_t \overset{\text{homot.}}{\simeq} \bigvee_{i=1}^{n-1} (\vee_{\mu_{t,i}} \boldsymbol{S}^i),$$

where $\mu_{t,i} \geq 0$ is the number of spheres of dimension i, with $\mu_{t,i} \geq \mu_{0,i}$, for any $i = 1, \ldots, n-1$;

(ii) *there exists an integrable vector field over M_t, null over P_t, which defines a continuous map $\tilde{\xi}_t : \partial M_t \to P_t$;*

(iii) *the fibre M_t is the mapping cylindre of $\tilde{\xi}_t$, and therefore M_t deformation retracts to P_t.*

Proposition 9. *For any $t \in \boldsymbol{D}^*_\eta$, there exist a continuous map $\Psi_t : M_t \to M_0$ and polyhedra $\mathcal{P}_t$ in M_t and $\mathcal{P}_0$ in M_0, both of (real) dimension $2n-3$, such that:*

(i) *$\mathcal{P}_t$ contains P_t and $\mathcal{P}_0$ contains P_0;*

(ii) *Ψ_t restricts to a homeomorphism $M_t \backslash \mathcal{P}_t \to M_0 \backslash \mathcal{P}_0$;*

(iii) *Ψ_t takes $\mathcal{P}_t$ to $\mathcal{P}_0$ and P_t to P_0;*

(iv) *When $n = 2$, we can consider $\mathcal{P}_t = P_t$ and $\mathcal{P}_0 = P_0$.*

We call Ψ_t a collapsing map for f.

To prove these propositions, we need to construct a good projection of the Milnor fibre X_t of f to a disk, denoted by $\varphi_t : X_t \to D_t$. This is what we do in the rest of this section:

For any linear form

$$l : \mathbb{C}^{n+1} \to \mathbb{C}$$

taking $0 \in \mathbb{C}^{n+1}$ to $0 \in \mathbb{C}$, the restriction of both f and l to M induces an analytic morphism

$$\phi_l : M \to \mathbb{C}^2$$

defined by $\phi_l(z) = \big(l(z), f(z)\big)$, for any $z \in M$

Lemma 10. *There exists a non-empty Zariski open set Ω in the space of non-zero linear forms of $\mathbb{C}^{n+1}$ to $\mathbb{C}$ that take $0 \in \mathbb{C}^{n+1}$ to $0 \in \mathbb{C}$, such that for any $l \in \Omega$, the analytic morphism $\phi_l : M \to \mathbb{C}^2$ satisfies:*

- (i) *if C is the critical set of ϕ_l and $\Gamma_l \subset M$ is the union of the irreducible components of C which are not contained in $f^{-1}(0)$, then Γ_l is either empty or a reduced complex curve;*
- (ii) *If $\Gamma_l \cap f^{-1}(0) = \{p_1, \ldots, p_r\}$, then for each p_i there exists a small neighbourhood V_i of p_i in M such that for any $x \in (\Gamma_l \cap V_i)\backslash\{p_i\}$, the hypersurface $\phi_l^{-1}(l(x), f(x))$ has an ordinary quadratic singularity at x;*
- (iii) *the restriction of ϕ_l to $\Gamma_l \cap V_i$, for each $i = 1, \ldots, r$, is finite and it defines an analytic morphism $(1-1)$ of $\Gamma_l \cap V_i$ to its image $\Delta_{l,i} := \phi(\Gamma_l \cap V_i)$;*
- (iv) *the tangent cone of $\Delta_{l,i}$ at $\phi_l(p_i)$ is $\mathbb{C} \times \{0\}$.*

If $l \in \Omega$, we say that l is a *general linear form* in respect to f, and we say that the curve Γ_l is the *polar curve of f* related to l and that the curve Δ_l is the *polar image of f* related to l.

From now on, we shall fix a good linear form l, and in order to simplify notation, we shall denote $\Omega := \Omega_l$, $\Gamma := \Gamma_l$ and $\Delta := \Delta_l$.

If we choose $\eta > 0$ sufficiently small, we can suppose that $\Gamma' := \Gamma \cap M_\eta$ is contained in the union $\cup_{i=1}^r V_i$. For a moment, let us denote by f' the restriction of f to M_η, and consider ϕ' defined on M_η by setting $\phi'(z) := \big(l(z), f'(z)\big)$. Without lost of generality, we can also suppose that $l(p_i) \neq l(p_j)$, for any $i \neq j$, $i, j \in \{1, \ldots, r\}$, so the previous lemma implies that the restriction of ϕ' to Γ' is finite and it defines an analytic morphism $(1-1)$ of Γ' to its image $\Delta' := \phi(\Gamma')$. In fact, if η is sufficiently small, then Γ' has exactly r-connected components $\Gamma'(p_i)$, which are curves in p_i, that is, Γ' is the disjoint union

$$\Gamma' = \bigsqcup_{i=1}^{r} \Gamma'(p_i),$$

and Δ' is the disjoint union

$$\Delta' = \bigsqcup_{i=1}^{r} \phi(\Gamma'(p_i)) := \bigsqcup_{i=1}^{r} \Delta'(p_i).$$

In order to simplify notation, from now on we shall denote $f := f'$ (defined on M_η), $\Gamma := \Gamma'$ and so on. See figure 1.



FIGURE 1.

Now fix $l \in \Omega$ such that the set $\mathcal{D} := l(M_\eta)$ is diffeomorphic to a disk in $\mathbb{C}$, and for any $t \in \mathbf{D}_\eta$, set

$$D_t := \mathcal{D} \times \{t\}.$$

Then $\phi : M_\eta \to \mathcal{D} \times \mathbf{D}_\eta$ induces a projection

$$\varphi_t : M_t \to D_t,$$

which is a fibre bundle over $D_t \backslash (\Delta \cap D_t)$.

5. PROOF OF PROPOSITIONS 8 AND 9 WHEN $n = 2$

Suppose that $M \subset \mathbb{C}^3$ has complex dimension 2. Then following the steps of [7] we construct a Lê Polyhedron and a collapse polyhedron for f as follows:

5.1. **The construction of the Lê Polyhedron.**

We have seen that, for each $t \in \mathbf{D}_\eta$,

$$\varphi_{t|} : M_t \backslash \varphi_t^{-1}(\Delta \cap D_t) \to D_t \backslash (\Delta \cap D_t)$$

is a topological fibre bundle, and therefore it is a topological covering of degree m_t.

Let $y_1(t), \ldots, y_k(t)$ be the points of the intersection $\Delta \cap D_t$. Note that each $y_j(t)$, for $j = 1, \ldots, k$, is contained in some $\Delta(p_i)$, for some $i = 1, \ldots, r$.

Let λ_t be the barycenter of the set of points $\{y_1(t), \ldots, y_k(t)\}$ in D_t and for each $j = 1, \ldots, k$, let $\delta(y_j(t))$ be a simple path (differentiable and with no double points) starting at λ_t and ending at $y_j(t)$, such that two of them intersect only at λ_t.

Set

$$Q_t := \bigcup_{j=1}^{k} \delta(y_j(t))$$

and

$$P_t := \varphi_t^{-1}(Q_t).$$

See figure 2.

Now, let v_t be a vector field in D_t such that v_t is:

- C^∞;
- null over Q_t;
- transversal to ∂D_t and points inwards.

Then the associated flow $q_t : [0,\infty[\times (D_t \backslash Q_t) \to D_t$ defines a map

$$\begin{array}{cccc} \xi_t & : & \partial D_t & \longrightarrow & Q_t \\ & & u & \longmapsto & \lim\limits_{\tau\to\infty} q_t(\tau,u) \end{array},$$

such that ξ_t is continuous, surjective and differentiable.

Since φ_t is a covering over $D_t \backslash Q_t$, which is differentiable in this case of dimension $n = 2$, we can lift v_t to a vector field E_t in M_t such that E_t is:

- continuous over M_t;
- differentiable over $M_t \backslash P_t$;
- null over P_t;
- integrable;
- transversal to ∂M_t and points inwards.

Then the associated flow $\tilde{q}_t : [0,\infty[\times (M_t \backslash P_t) \to M_t$ defines a map

$$\begin{array}{cccc} \tilde{\xi}_t & : & \partial M_t & \longrightarrow & P_t \\ & & z & \longmapsto & \lim\limits_{\tau\to\infty} \tilde{q}_t(\tau,z) \end{array},$$

such that $\tilde{\xi}_t$ is continuous, surjective and differentiable.

So now we have to show that M_t is homeomorphic to the mapping cylinder of $\tilde{\xi}_t$. In fact, the integration of the vector field $\tilde{v}_t$ gives a surjective continuous map

$$\alpha : [0,\infty] \ \times \ \partial M_t \to M_t$$

that restricts to a diffeomorphism

$$\alpha_| : [0,\infty[\times \partial M_t \to M_t \backslash P_t.$$

Since the restriction $\alpha_\infty : \{\infty\} \times \partial M_t \to P_t$ is equal to $\tilde{\xi}_t$, which is differentiable and surjective, it follows that the induced map

$$[\alpha_\infty] : \big((\{\infty\} \times \partial M_t)/ \sim \big) \to P_t$$

is a homeomorphism, where $\sim$ is the equivalence relation given by the identification $(\infty, z) \sim (\infty, z')$ if $\alpha_\infty(z) = \alpha_\infty(z')$. Hence the map

$$[\alpha] : \big(([0,\infty] \times \partial M_t)/ \sim \big) \to M_t$$

induced by α defines a homeomorphism between M_t and the mapping cylinder of $\tilde{\xi}_t$.

This proves proposition 8 for the case $n = 2$. The following proposition will be useful in the last section of this paper.



FIGURE 2.

Proposition 11. *For any* $t \in \boldsymbol{D}_\eta^*$*, the topology of* P_t *is described by the pair* (m_t, k)*, that is, two polyhedra* $P_t(1)$ *and* $P_t(2)$ *construct as above are homeomorphic if, and only if,* $m_t(1) = m_t(2)$ *and* $k(1) = k(2)$*. In the same way, the topology of* P_0 *is described by the pair* (m_0, r)*.*

Proof. Note that:

- $\varphi_t^{-1}(y_j(t))$ is contained in Γ. In fact, since the restriction $\varphi_{t|} : M_t \backslash (\Gamma \cap M_t) \to Im(\varphi_{t|}) \subset D_t$ is a surjective submersion, it follows from Ehresmann's fibration lemma that it is a topological fibre bundle. Then, if there exists $z \in M_t \backslash (\Gamma \cap M_t)$ such that $\varphi_{t|}(z) = y_j(t)$, it follows that $y_j(t) \in Im(\varphi_{t|})$ and then $\varphi_{t|}^{-1}(y_j(t))$ would be given by m_t points in $M_t \backslash (\Gamma \cap M_t)$, and hence $\varphi_t^{-1}(y_j(t))$ would be given by $(m_t + a)$ points in M_t, for some $a \geq 1$. But then the Lê Polyhedron would not be connected, since there are just m_t paths connecting $\varphi_t^{-1}(y_j(t))$ and $\varphi_t^{-1}(\lambda_t)$. This is a contradiction since we have already proved that the connected manifold M_t deformation retracts to P_t;
- Therefore $\varphi_t^{-1}(y_j(t))$ is just one point, by item (iii) of lemma 10;
- Hence the topology of P_t is encoded in the pair (m_t, k). In fact, for any $t \in \mathbf{D}_\eta^*$, it is easy to see that P_t has exactly $(m_t + k)$ vertices and it has the homotopy type of a bouquet of $(m_t - 1)(k - 1)$ circles if $m_t > 1$ and $k > 1$, or that P_t is contractible if $m_t = 1$ or $k = 1$;
- Also, one can check that P_0 has exactly $(m_0 + r)$ vertices and it has the homotopy type of a bouquet of $(m_0 - 1)(r - 1)$ circles if $m_0 > 1$ and $r > 1$, or that P_0 is contractible if $m_0 = 1$ or $r = 1$.

□

5.2. **The collapse along a path.** We can do the construction of the vector field E_t simultaneously for all t in a simple path γ in $\mathbf{D}_\eta$ joining 0 and some $t_0 \in \partial \mathbf{D}_\eta$, such that γ is transverse to $\partial \mathbf{D}_\eta$. To simplify, we shall assume that γ is the closed segment of line in $\mathbf{D}_\eta$ joining 0 and t_0.

The natural projection $\pi : \mathcal{D} \times \mathbf{D}_\eta \to \mathbf{D}_\eta$ restricted to Δ induces a ramified covering

$$\pi_| : \Delta \to \mathbf{D}_\eta$$

whose ramification locus is $D_0 \cap \Delta = \{\phi(p_1), \ldots, \phi(p_r)\}$, by item (iv) of lemma 10.

Hence the inverse image of $\gamma \backslash \{0\}$ by this covering defines k disjoint simple paths in Δ, and each one of them is diffeomorphic to $\gamma \backslash \{0\}$. Each of these paths have $\phi(p_i)$ in its closure, for some $i = 1, \ldots, r$, and it contains the points $y_j(t)$, for some $j = 1, \ldots, k$ and any $t \in \gamma \backslash \{0\}$. We shall denote by $\varsigma_{i,j}$ the respective path that has $\phi(p_i)$ in its closure and contains $y_j(t)$. In particular, we have that $r \leq k$. See figure 3.

Moreover, the set $\Lambda = \bigcup_{t \in \gamma} \lambda_t$ defines a simple path in $\mathcal{D} \times \mathbf{D}_\eta$ such that either $\Lambda \cap \Delta = \phi(p_1)$, if $r = 1$, or $\Lambda \cap \Delta = \emptyset$, if $r > 1$.

We can choose the paths $\delta(y_j(t))$ in such a way that

$$T_j := \bigcup_{t \in \gamma} \delta(y_j(t))$$

forms either a triangle, if $r = 1$, or a square, if $r > 1$, differentiably immersed in

$$\bigcup_{t \in \gamma} D_t = \mathcal{D} \times \gamma$$

outside $\delta(y_j(0))$. For any $j, j' \in \{1, \ldots, k\}$ with $j \neq j'$, note that either $T_j \cap T_{j'} = \Lambda$, if both $\varsigma_{i,j}$ and $\varsigma_{i',j'}$ are defined for some $i, i' \in \{1, \ldots, r\}$ with $i \neq i'$; or $T_j \cap T_{j'} = \Lambda \cup \gamma(y_j(0)) = \Lambda \cup \gamma(y_{j'}(0))$, if both $\varsigma_{i,j}$ and $\varsigma_{i,j'}$ are defined for some $i \in \{1, \ldots, r\}$. See figure 4. Set

$$Q := \bigcup_{j=1}^{k} T_j.$$



Figure 3.



Figure 4.

Now, let V be a vector field in $\mathcal{D}\times\gamma$ such that V is:

- continuous;
- null over Q;
- differentiable over $(\mathcal{D}\times\gamma)\backslash Q$;
- transversal to $\partial\mathcal{D}\times\gamma$.

Then the associated flow $w : [0,\infty[\, \times \big((\mathcal{D}\times\gamma)\backslash Q\big) \to \mathcal{D}\times\gamma$ defines a map

$$\begin{array}{cccc} \xi & : & \partial\mathcal{D}\times\gamma & \longrightarrow & Q \\ & & z & \longmapsto & \lim\limits_{\tau\to\infty} w(\tau,z) \end{array},$$

such that ξ is continuous, surjective and differentiable.

For any real $A>0$, set

$$V_A(Q) := (\mathcal{D}\times\gamma)\backslash w\big([0,A[\times\partial\mathcal{D}\times\gamma\big),$$

a closed neighbourhood of Q in $\mathcal{D}\times\gamma$. Note that $\partial V_A(Q)$ is a differentiable manifold that fibres over γ with fibre a circle, by the restriction of the projection π. Moreover, $\mathcal{D}\times\gamma$ is clearly the mapping cylinder of ξ.

Since

$$\phi_| : M_\eta\backslash\phi^{-1}(Q) \to (\mathcal{D}\times\mathbf{D}_\eta)\backslash Q$$

is a locally trivial fibration, it follows that $\phi^{-1}(\partial V_A(Q))$ is a differentiable submanifold of M_η which is a fibre bundle over γ.

Now, set

$$P_\gamma := \phi^{-1}(Q),$$

which we call the *collapse polyhedron of f along* γ. It is a polyhedron in M_η of real dimension n. Let θ be a vector field in γ that goes from t_0 to 0 in time $a>0$.

Set

$$Z := \big(M_\eta\cap f^{-1}(\gamma)\big)\backslash P_\gamma.$$

Since

$$Z = \phi^{-1}\big((\mathcal{D}\times\gamma)\backslash Q\big) \xrightarrow{\phi} (\mathcal{D}\times\gamma)\backslash Q \xrightarrow{\pi} \gamma$$

and

$$\phi^{-1}\big(\partial V_A(Q)\big) \xrightarrow{\phi} \partial V_A(Q) \xrightarrow{\pi} \gamma$$

are differential fibre bundles, we can lift θ to obtain a vector field E such that:

- E is differentiable;
- E is tangent to $\phi^{-1}\big(\partial V_A(Q)\big)$, for any $A > 0$.

Then the associated flow $g : [0,a] \times Z \to Z$ defines a C^∞-diffeomorphism Ψ from $M_{t_0} \backslash P_{t_0}$ to $M_0 \backslash P_0$ that extends to a continuous map from M_{t_0} to M_0 and that sends P_{t_0} to P_0.

This proves proposition 9 for the case $n = 2$.

Remark 12. Note that since $M_{t_0} \backslash P_{t_0}$ is diffeomorphic to $M_0 \backslash P_0$, we must have that $m_0 = m_t$. Also note that, for any path γ, we have that P_γ deformation retracts to P_0.

5.3. **The collapse along a disk.** We can go further and describe the collapse simultaneously along all the disk $\mathbf{D}_\eta$ in the following way:

Consider the intersection of Δ with $\mathcal{D} \times \mathbf{D}^*_\eta$. Then we obtain k punctured disks Υ_j. The barycenter points of these punctured disks also give a punctured disk Λ such that the intersection of the closure of all this punctured disks is either $\phi(p_1)$, if $r = 1$, or empty, if $r > 1$. Fix $j \in \{1, \ldots, k\}$. The union of paths $\delta(y_j(t))$ for all $t \in \mathbf{D}_\eta$ gives a 3-dimensional polyhedron T_j in $\mathcal{D} \times \mathbf{D}_\eta$. Note that either $T_j \cap T_{j'} = \Lambda$ or $T_j \cap T_{j'} = \Lambda \cup \gamma(y_j(0)) = \Lambda \cup \gamma(y_{j'}(0))$.

Then we define

$$Q := \bigcup_{j=1}^{k} T_j$$

and, as before, we construct a vector field V in $\mathcal{D} \times \mathbf{D}_\eta$ that retracts $\mathcal{D} \times \mathbf{D}_\eta$ onto Q. Now set

$$P := \phi^{-1}(Q),$$

which we call the *collapse polyhedron of f along a disk.* It is a polyhedron of real dimension $n+1$, contained in M_η.

Also set

$$Z := M_\eta \backslash P.$$

Since ϕ is a submersion over $(\mathcal{D} \times \mathbf{D}_\eta) \backslash Q$, it follows that V lifts to a vector field E in Z with the desired properties, as before. In particular, we have:

Corollary 13. *M_η deformation retracts to P.*

---

6. Proof of propositions 8 and 9 when $n \geq 3$

In this section we prove propositions 8 and 9 when M has dimension $n \geq 3$. We do it by induction on n, since we already have the lemmata for $n = 2$. Suppose the results are true for $n - 1$.

6.1. **The construction of the Lê Polyhedron.** We already know that, for each $t \in \mathbf{D}_\eta$, the projection

$$\varphi_t : M_t \to D_t$$

determines a topological fibre bundle

$$\varphi_{t|} : M_t \backslash \varphi_t^{-1}\big(\{y_1(t), \ldots, y_k(t)\}\big) \to D_t \backslash \{y_1(t), \ldots, y_k(t)\},$$

where $\{y_1(t), \ldots, y_k(t)\} = \Delta \cap D_t$. Note that each $y_j(t)$, for $j \in \{1, \ldots, k\}$, is contained in some $\Delta(p_i)$, $i \in \{1, \ldots, r\}$, as defined in section 4, and that $k \geq r$ if $t \neq 0$, and $k = r$ if $t = 0$.

Let u' be a point in $\mathcal{D}$ such that the complex line $\{u'\} \times \mathbb{C}$ is not a component of Δ. If we set $\lambda_t := (u', t)$, then $\lambda_t \notin \{y_1(t), \ldots, y_k(t)\}$. For each $j = 1, \ldots, k$, let $\delta(y_j(t))$ be a simple path starting at λ_t and ending at $y_j(t)$, such that two of them intersect only at λ_t.

Now, first we consider f' the restriction of f to $M_\eta \cap \{l = u'\}$, which we can suppose is a differentiable complex manifold of dimension $n-1$, and then we can apply proposition 8, by induction hypothesis, to obtain a Le Polyhedron P'_t and a vector field E'_t as in the previous section. Second, recall that ϕ defines a fibre bundle over $\delta(y_j(t)) \backslash \{y_j(t)\}$.

Finally, for each $j = 1, \ldots, k$, define the point

$$x_j(t) := \phi^{-1}\big(y_j(t)\big) \cap \Gamma,$$

which is the isolated singularity of $\phi^{-1}(y_j(t))$. If we look at the local situation at $x_j(t)$, we can use the result of [7], that is, we can consider B_j a small ball in $\mathbb{C}^{n+1}$ centered at $x_j(t)$ (contained in M_η) and D_s a small disk in D_t centered at $y_j(t)$ such that the restriction

$$\varphi_{t|} : B_j \cap \varphi_t^{-1}(D_s) \to D_s$$

satisfies the hypothesis (2.3.2) and (2.3.3) of [7]. See figure 5.



FIGURE 5.

Then if D_s^+ denotes a semi-disk in D_t containing $D_s \cap \delta(y_j(t))$, we obtain a collapse cone P_j and a vector field E_j in $\varphi_t^{-1}(D_s^+) \cap B_j$. Set

$$a_j := \partial D_s \cap \delta(y_j(t)).$$

Then

$$P_j(a_j) := \varphi_t^{-1}(a_j) \cap P_j$$

is a Lê Polyhedron for the germ $\varphi_t : (M_t, x_j(t)) \to (D_t, y_j(t))$.

Next we construct some useful vector fields on $A_j := \varphi_t^{-1}\big(\delta(y_j(t))\backslash\{y_j(t)\}\big)$:

- **Vector Field Ξ:** let ξ be a C^∞ vector field non-zero on $\delta(y_j(t))\backslash\{y_j(t)\}$ that goes from $y_j(t)$ to $u' = \lambda_t$. Since φ_t is a fibre bundle over $\delta(y_j(t))\backslash\{y_j(t)\}$, we can lift ξ to a vector field Ξ on A_j which is integrable (see [13] and (2.3.2.2) of [7]) and tangent to $\mathring{A}_j$ and ∂A_j;
- **Vector Field $\mathcal{V}$:** We can transport the vector field E'_t of $\varphi_t^{-1}(u')$ to all the fibres $\varphi_t^{-1}(u)$, for any $u \in \delta(y_j(t))\backslash\{y_j(t)\}$. Then we obtain a vector field $\mathcal{V}$ on A_j whose restriction to $\varphi_t^{-1}(u')$ is E'_t and the restrictions to $\varphi_t^{-1}(u)$ are the vector fields of $\varphi_t^{-1}(u)$;
- **Vector Field Θ:** Let θ be a differentiable function on $\delta(y_j(t))$ such that $\theta(u') = 0$ and such that θ is non-zero and positive on $\delta(y_j(t))\backslash\{u', y_j(t)\}$. Let Θ be the vector field on A_j defined by θ;
- **Vector Field $\mathcal{V}_1$:** Define $\mathcal{V}_1 := \mathcal{V} + \Theta\Xi$, which is integrable, tangent on the interior of A_j and transverse and pointing inwards on the boundary ∂A_j.

Since $\mathcal{V}$ and Ξ are transversal, the vector field $\mathcal{V}_1$ is zero only on the Lê Polyhedron P'_t of $\varphi_t^{-1}(u')$. Then if z is a point in $A_j\backslash\varphi_t^{-1}(u')$, the orbit of $\mathcal{V}_1$ that passes through z has its limit point z'_1 in P'_t.

Moreover, since the orbit of $\mathcal{V}$ that passes through z has its limit point z' in the transportation of P'_t to $\varphi_t^{-1}\big(\varphi_t(z)\big)$ by Ξ, it follows that z'_1 is the point corresponding to z' by Ξ. That comes from the fact that $\mathcal{V}$ and Ξ commute by construction.

Now, since $P_j(a_j)$ is obviously contained in A_j, it follows that $\mathcal{V}_1$ takes $P_j(a_j)$ to P'_t. In fact, it takes all the fibre $\varphi_t^{-1}(a_i) \cap B_r$ to P'_t. Now, since the action of the flow given by $\mathcal{V}$ is simplicial, we can assume that the action of the flow given by $\mathcal{V}_1$ is simplicial. Then the image of $P_j(a_j)$ by the action of $\mathcal{V}_1$ is a sub-polyhedron P'_j of P'_t. Moreover, the orbits of the points in $P_j(a_j)$ give a polyhedron R_j.

Set

$$\tilde{P}_j := P_j \cap \varphi_t^{-1}\big(\delta(y_j(t))\big)$$

and

$$S_j := \tilde{P}_j \cup R_j \cup P'_j.$$

See figure 6.

Lemma 14. *Let $r > 0$ be small enough such that the ball $B_r(x_j(t))$ of radius r centered at $x_j(t)$ is contained in B_j, for any $j = 1, \ldots, k$, and then redefine $B_j := B_r(x_j(t))$. Then, for any $j_1, j_2 = 1, \ldots, k$, we have:*

(i) $P'_{j_1} = P'_{j_2}$*;*

(ii) $P_{j_1}(a_{j_1}) \overset{\text{diff.}}{=} P_{j_2}(a_{j_2})$*;*

(iii) $S_{j_1} \overset{\text{diff.}}{=} S_{j_2}$*;*

(iv) $S_j \overset{\text{diff.}}{=} S_i^0$*, for any* $j = 1, \ldots, k$ *and* $i = 1, \ldots, r$*.*



Figure 6.

Proof. The natural extension of Ξ to $[\delta(y_{j_1}(t))\backslash\{y_{j_1}(t)\}] \cup [\delta(y_{j_2}(t))\backslash\{y_{j_2}(t)\}]$ gives (i) and (ii). Clearly, $R_j \overset{\text{diff.}}{=} P'_j \times I$, for any $j = 1, \dots, k$; where I denotes a real interval, and then (iii) follows. Finally, since $\phi_| : \tilde{M}_\eta \backslash \phi^{-1}(\Delta) \to (\mathcal{D} \times \mathbf{D}_\eta)\backslash\Delta$ is a fibre bundle, it follows that $\varphi_t^{-1}(u) \overset{\text{diff.}}{=} \varphi_0^{-1}(u)$, for any u in $D_t \backslash (\Delta \cap D_t)$, and then

$$P'_t \overset{\text{diff.}}{=} P'_0.$$

But this implies that $P'_j \overset{\text{diff.}}{=} P'^0_i$, for any $j = 1, \dots, k$ and $i = 1, \dots, r$, where P'^0_i is the sub-polyhedron of P'_0 defined analogous to the way P'_j is defined. Hence $S_j \overset{\text{diff.}}{=} S^0_i$.

□

Then the polyhedron in M_t we are looking for is given by

$$P_t := P'_t \bigcup_{j=1}^{k} S_j$$

and the special polyhedron in M_0 is given by

$$P_0 := P'_t \bigcup_{j=1}^{r} S_j.$$

Figure 7 illustrates the homotopy type of both P_t and P_0, as well as the degeneration of the first to the second one, which wel construct next.

Once the Lê Polyhedrons P_t and P_0 are defined, the proof of proposition 8 follow the very same arguments of sections (4.2.1), (5.1) and (5.2) of [7].

6.2. **The collapsing map.** Consider the disk

$$\Lambda := \{u'\} \times \mathbf{D}_\eta.$$



FIGURE 7.

Then we define T_j for $j = 1, \dots, k$ as in 5.3, that is,

$$T_j := \bigcup_{t \in \mathbf{D}_\eta} \delta(y_j(t)),$$

and the sets

$$Q_t := \bigcup_{j=1}^{k} \delta(y_j(t)),$$

$$Q_0 := \bigcup_{i=1}^{r} \delta(\phi(p_i))$$

and

$$Q := \bigcup_{j=1}^{k} T_j = \bigcup_{t \in \mathbf{D}_\eta} Q_t = \bigcup_{\substack{t \in \mathbf{D}_\eta \\ 1 \le j \le k}} \delta(y_j(t)).$$

Then we consider $\mathcal{V}$ a differentiable vector field in $\mathcal{D} \times \mathbf{D}_\eta$ which is transversal to each $D_t = \mathcal{D} \times \{t\}$, points inwards, is non-null in $\mathcal{D} \times \mathbf{D}^*_\eta$, and is tangent to Q and to Λ.

Hence the lift $\tilde{\mathcal{V}}$ of $\mathcal{V}$ by ϕ determines a vector field in M_η, which gives the collapse polyhedron P as the union of the orbits passing through the points of all the polyhedra P_t's, for $t \in \mathbf{D}^*_\eta$. Moreover, $\tilde{\mathcal{V}}$ defines the (continuous) collapsing map $\Psi_t : M_t \to M_0$, for each $t \in \mathbf{D}^*_\eta$ with the desired properties, setting

$$\mathcal{P}_t := \phi^{-1}(Q_t)$$

and

$$\mathcal{P}_0 := \phi^{-1}(Q_0).$$

Then we have proved proposition 9. In fact, (i) and (iii) are obvious and (ii) follows from the fact that the restriction $\phi_| : M_\eta \backslash \phi^{-1}(Q) \to (\mathcal{D} \times \mathbf{D}_\eta) \backslash Q$ is a topological fibre bundle. Finally, (iv) is clear by the definition of the Lê Polyhedron when $n = 2$. We also have proved:

Corollary 15. *M_η deformation retracts to P.*

Remark 16. *Note that P_t is homeomorphic to P_0 if, and only if, $r = k$; which happens if, and only if, $\mathcal{P}_t$ is homeomorphic to $\mathcal{P}_0$; which happens if, and only if, M_t is homeomorphic to M_0.*

---

7. Proof of lemmas 4, 5, 6 and 7

Lemma 4 is follows directly applying proposition 8 to the projection π_t and observing that $P_{t,0}$ has the homotopy type of a bouquet of spheres of dimension $n-1$, since $(\{0\} \times \mathbf{B}_{\epsilon_2})$ is a Milnor ball for f_0, and hence, by lemma 14, $P_{t,s}$ has the homotopy type of a bouquet of spheres of dimension $n-1$, for any $s \in \Sigma_\epsilon$.

Lemma 6 follows setting P_t to be the collapse polyhedron P of π_t along Σ_ϵ, since we have showed that $M_t = X_t$ deformation retracts to P (see corollary 15).

Now, since the construction of the Lê Polyhedron of both π_t and f_s depends only on the projection φ_t, which is the restriction of the linear form l to the Milnor fibre of the function, lemma 7 is trivial: when constructing a Lê Polyhedron for f_s, one just have to consider the linear form l to be the same of that used in the construction of $P_{t,s}$. Note that since the set of general linear forms l, denoted by Ω, is a non-empty Zariski set in the space of all the linear forms of $\mathbb{C}^n$ to $\mathbb{C}$, we can assume that l is a good one (otherwise, just take another linear form l in the construction of $P_{t,s}$).

Finally, lemma 5 follows from proposition 9, observing that:

- Since $P_{t,0}$ is a Lê Polyhedron for the isolated singularity f_0 inside a Milnor ball, it follows from the main theorem of [7] that we can consider $\mathcal{P}_{t,0} = P_{t,0}$ and $P_{0,0} = \{0\}$;
- Since $\tilde{\Psi}$ takes $\mathcal{P}_{t,s}$ to $\mathcal{P}_{t,0}$ and $P_{t,s}$ to $P_{t,0}$, it follows that $\mathcal{P}_{t,s} = P_{t,s}$, for any $s \in \Sigma_\epsilon$ (in fact, $\mathcal{P}_{t,s} = (\tilde{\Psi}_{t,s})^{-1}(\mathcal{P}_{t,0}) = (\tilde{\Psi}_{t,s})^{-1}(P_{t,0}) = P_{t,s}$).

---

8. The μ-constant problem

Two germs of complex analytic hypersurfaces at the origin in $\mathbb{C}^n$, denoted by $(X_1, 0)$ and $(X_2, 0)$, are said to be topologically equisingular (or to have the same embedded topological type) if there exists a local homeomorphism $h : (\mathbb{C}^n, 0) \to (\mathbb{C}^n, 0)$ such that $h(X_1) = X_2$.

Definition 17. *Let $f_s : (\mathbb{C}^n, 0) \to (\mathbb{C}, 0)$ be a family of holomorphic isolated singularity germs depending holomorphically on the parameter $s \in \mathbb{C}$. It defines a family of germs of isolated singularity hypersurfaces $X_s := \big(f_s^{-1}(0), 0\big)$, which is said to be embedded topologically equisingular (or to have constant embedded topological type) along $\mathbb{C} \times \{0\}$ if, for any s sufficiently small, X_s and X_0 have the same embedded topological type at the origin.*

In this section, we use the construction of a Lê Polyhedron for line singularities to prove that the family X_s as above has constant embedded topological type if it admits an uniform Milnor radius and the Milnor number $\mu(f_s)$ is independent of s.

Consider

$$f : (\mathbb{C}^{n+1}, 0) \to (\mathbb{C}, 0)$$

the holomorphic function defined by the family f_s, that is, $f(z_1, \ldots, z_{n+1}) = f_{z_{n+1}}(z_1, \ldots, z_n)$. The singular set of f is clearly the z_n-axis. Also consider the hyperplane sections $H_s = \{z_{n+1} = s\}$, for each $s \in \Sigma$. As in section 3, define the Milnor fibration of f using polydiscs, that is, if ϵ_1 and ϵ_2 are sufficiently small positive reals with $0 < \epsilon_1 < \epsilon_2 << 1$, one has that the restriction

$$f_| : f^{-1}(\mathbf{D}^*_\eta) \cap (\mathbf{B}_{\epsilon_1} \times \mathbf{B}_{\epsilon_2}) \to \mathbf{D}^*_\eta$$

is a fibre bundle, where $\mathbf{B}_{\epsilon_1}$ denotes the ball in $\Sigma = \mathbb{C}$ and $\mathbf{B}_{\epsilon_2}$ denotes the ball in $H_s = \mathbb{C}^n$.

Now suppose that f_s admits an uniform Milnor radius. Then we can take $\epsilon > 0$ sufficiently small such that the ball $\mathbf{B}_\epsilon(s) = \mathbf{B}_\epsilon \cap H_s$ is a Milnor ball for f_s, for any $s \in \Sigma$. Fix $t \in \mathbf{D}^*_\eta$ and for each $s \in \Sigma$, consider the projection $\pi_t : X_t \to \Sigma$ and the polyhedron $P_{t,s}$ in $f_s^{-1}(t) \cap \mathbf{B}_\epsilon$ as constructed in the previous sections, which is a Lê Polyhedron for the isolated singularity germ f_s, and hence it has the homotopy type of a bouquet of μ-spheres of dimension $n-1$.

We want to show that $P_{t,s}$ is homeomorphic to $P_{t,0}$, and hence Lemma 5 implies that $\pi_t^{-1}(s)$ is homeomorphic to $\pi_t^{-1}(0)$, that is, $f_s^{-1}(t) \cap \mathbf{B}_\epsilon$ is homeomorphic to $f_0^{-1}(t) \cap \mathbf{B}_\epsilon$. When $n - 2$, Proposition 11 implies that

$$(m_t - 1)(r - 1) = \mu = (m_t - 1)(k - 1),$$

and hence remark 12 implies $r = k$. Then the pairs (m_t, k) and (m_0, r) are equal, and therefore $P_{t,s}$ is homeomorphic to $P_{t,0}$.

When $n \geq 3$, by the construction of the Lê Polyhedron we have made in section 5.1 we know that $P_{t,s}$ is the union of k-copies of a polyhedron S, glued together on a sub-polyhedron P of S, and that $P_{t,0}$ is the union of r-copies of S, glues together on P. Now, since $P_{t,s}$ is homotopy equivalent to $P_{t,0}$, we must have $k = r$, and therefore $P_{t,s}$ is homeomorphic to $P_{t,0}$.

So we have proved that if $f_s : (\mathbb{C}^n, 0) \to (\mathbb{C}, 0)$ is a family of germs of holomorphic isolated singularities depending holomorphically on the parameter $s \in \mathbb{C}$ which admits an uniform Milnor radius and is μ-constant, then the embedded topological type of of the Milnor fibre of f_s does not depend on s.

Then obviously the embedded topological type of the boundary of the Milnor fibre of f_s does not depend on s. Since f_s is an isolated singularity, the boundary of its Milnor fibre has the same embedded topological type of the embedded link of X_s (that is, the homeomorphism class of the pair $\big(\mathbf{S}_\epsilon, \mathbf{S}_\epsilon \cap X_s\big)$, which is independent of ϵ if ϵ is small enough - see [5]).

Since the embedded link of X_s is constant along the family, it follows from the conical structure of singularities that the family has constant embedded topological type. Then we have proved:

Theorem 18. *Let $f_s : (\mathbb{C}^n, 0) \to (\mathbb{C}, 0)$ be a family of complex isolated singularity map-germs depending holomorphically on the parameter $s \in \mathbb{C}$ which admits an uniform Milnor radius, and consider the family of germs of isolated singularity hypersurfaces X_s in $\mathbb{C}^n$ defined by f_s. Then X_s has constant embedded topological type if, and only if, the Milnor number $\mu(f_s)$ is independent of s.*

References

[1] K. Bekka, *Regular Quasi-Homogeneous Stratifications*, "Stratification, singularities and diferential equations II Stratifications and Topology of Singular Space", Travaux en cours 55, Hermann, 1-14, 1997.

[2] K. Bekka, *(c)-Regularité et Trivialité Topologique*, Singularity theory and its applications, Warwick, 1989, Part I, D. Mond and J. Montaldi, Eds., SLNM 1462, Springer, (1991), pp.42-62.

[3] K. Bekka and S. Koike, *The Kuo Condition, an Inequality of Thoms Type and (c)-Regularity*, Topology, vol. 37, 1, pp.4562, 1998.

[4] J.F. de Bobadilla, *Answers to some equisingularity questions*, Invent. Math. 161 (2005), 657-675.

[5] A.H. Durfee, *Neighbourhoods of algebraic sets* Trans. Am. Math. Soc., 276 (1983), 517-530.

[6] A. Horváth and A. Némethi, *On the Milnor fiber of non-isolated singularities*, Studia Sci. Math. Hungar. 43 (2006), no. 1, 131136.

[7] D.T. Lê, *Polyèdres èvanescents et effondrements*, A fte of topology, 293329, Academic Press, Boston, MA, 1988.

[8] D.T. Lê, *Calcul du nombre de cycles èvanouissants d'une hypersurface complexe*, Ann. Inst. Fourier 23 (1973), 261-270, Grenoble, France.

[9] D.T. Lê and C.P. Ramanujam, *The invariance of Milnors number implies the invariance of the topological type*, Am. J. Math. 98, 67-78 (1976).

[10] J.W. Milnor, *Singular points of complex hypersurfaces*, Ann. of Math. Studies 61, Princeton, 1968.

[11] D. Siersma, *Isolated line singularities*, Singularities, Part 2 (Arcata, Calif., 1981), 485-496, Proc. Sympos. Pure Math., 40, Amer. Math. Soc., Providence, RI, 1983.

[12] E. Spanier, *Algebraic Topology*, McGraw-Hill, 1966.

[13] J. L. Verdier, *Stratifications de Whitney et théorèm* de Bertini-Sard, Inv. Math. 36 (1976), 295-312.

Instituto de Matemáticas, Unidad Cuernavaca, Universidad Nacional Autónoma de México

E-mail address: `lelobot@gmail.com`